CAMP EMERGENCY MEDICATION PLAN

Scout'sName:		_ Date of Birth:	Pack or Troop #:	
Cam	p Hinds Cam	p Telephone & Fax:	207-655-4878	
TO BE COMPLETED BY PARENT	OR GUARDIAN:			
I authorize the exchange of medical	information about my child	's asthma between the	physician's office and camp nurse.	
Parent or Guardian signature:				
Parent or Guardian tel.# home:	work:_		_ cell phone:	
Physician/Healthcare Provider Name:		Parent concerns:		
My child may carry and use his/her:	inhaled asthma medicine	□ Yes □ No	Epi-Pen □ Yes □ No □ N/	Α
TO BE COMPLETED BY CAMPER		•		
Provider name:			Fax#	
	□ NO changes	from previous pla	an	
Peak Flow:				
Child's predicted, or personal best				
Child's Green Zone:	Yellow Zone:		Red Zone: below	
Medications:				
	ications:			
Preventive (Controller) Medi	cations:			
Quick Relief Medications: (ca			• • •	
	,			
Inhaler with spacer OR nebulize	er Dose/Frequency:			
Allergies /Triggers for asthm	na: □ None known			
□ Avoid animals	.ia. = Nono knom			
☐ Other triggers to avoid:				
Exercise Pretreatment Instru				
			or	
			child's peak flow is below	
Asthma Exacerbation Treatr			•	
		** · · · · · · · · · · · · · · · · · ·		
➤ YELLOW ZONE: If child is cou		=		
☐ Give 2 puffs of child's quick reli recover to Green Zone. Notif	ief inhaler with spacer (or new v parents of exacerbation.	oulizer treatment). Iviay	be repeated in 10 minutes if doesn't	
Other:	y paromo or oxaconomic			
> RED ZONE: If child is in respi	irotory distress and/or pea	k flow is in Red 7one:		
☐ Give 4 puffs quick relief inhaler	•		care Provider	
Call 911 if child does not im				
☐ Other:				
Special Instructions:				
	o carrv and use inhaled me	dications and Epi-pen	after demonstrating appropriate use of	of
Inhalers and or Epi-Pen to camp n	urse. Please check approp	riate boxes below:	and domented appropriate	••
This camper has the knowledge	and skill to carry and use:	☐ Inhaled medicatio	n □ Epi-pen	
> This camper is not able to carry	and use by himself/herself:	☐ Inhaled medication	on 🛘 Epi-pen	
	-		nes more than 2 times a week (i.e. in exce	SS
Other:				
Healtho	care Provider signature	D	ate	
TO BE COMPLETED BY CAMP I	NURSE: This camper	demonstrates know	ledge and skill to carry and use:	
Inhaler medications	□ NO _			
Epi-Pen ☐ YES	□ NO □ N/A	Camp Nurse Sign	nature Date	

CAMP EMERGENCY MEDICATION PLAN

ASTHMA PLAN INSTRUCTIONS

Every camper with asthma should have a current

Camp Emergency Medication Plan completed and signed by their physician (or other health care provider) and kept on file in the camp nurse's office. The form must also be signed by a parent/guardian. The plan should be updated each year or when there are major changes to the plan (such as in medication type or dose). The physician's office is encouraged to fax the plan to the camper's camp nurse.

The camp plan is intended to strengthen the partnership of families, healthcare providers and the camp. It is based on the NHLBI Guidelines for Asthma Management. (For more information contact the camp nurse or www.nhlbi/nih/gov).

CARRYING AND ADMINISTERING QUICK RELIEF INHALERS and/or Epi-Pen:

- Most campers are capable of carrying and using their quick relief inhaler by themselves. The camper, camper's parents, camp nurse and healthcare provider should make this decision. The camp nurse must also evaluate technique for effective use.
- The appropriate boxes must be checked by the parent, provider and camp nurse to indicate the camper's ability to carry and self-administer these medications.

USE OF QUICK RELIEF MEDICATIONS MORE THAN TWICE WEEKLY:

> This indicates poor control of asthma, and providers should be notified by the camp nurse or designated staff.

PEAK FLOW ZONES (based on camper's personal or predicted best):

Green zone: Peak flow 80-100%

- ➤ Symptoms and/or use of quick relief medication < 2 times a week
- > Use daily controller medication at home
- > Full participation in physical education and sports

Yellow zone: Peak flow 50-80%

- ➤ Has symptoms or needs quick relief medication >2 times a week
- > Needs quick relief medication and further observation by camp nurse; notify parents
- > Attend physical education but restrict strenuous aerobic activity

Red zone: Peak flow <50%

- Symptoms may include shortness of breath, retractions, difficulty talking or walking; quick relief medication not effective
- > Requires immediate action, close monitoring and notification of parent and healthcare provider